




岡田忠司
大正十一年

昭和十一年八月二十一日

岡田忠司



δ (ppm)

ured at 25°C: (A) in CS₂ solution; (B) in

の混合液は市販品 (東京化成工業) を用いた。

(A)






Table 1. Molecular weight of the compounds.



Compound	Molecular weight
1	284.4
2	284.4
3	284.4
4	284.4
5	284.4
6	284.4
7	284.4
8	284.4
9	284.4
10	284.4
11	284.4
12	284.4
13	284.4
14	284.4
15	284.4
16	284.4
17	284.4
18	284.4
19	284.4
20	284.4
21	284.4
22	284.4
23	284.4
24	284.4
25	284.4
26	284.4
27	284.4
28	284.4
29	284.4
30	284.4
31	284.4
32	284.4
33	284.4
34	284.4
35	284.4
36	284.4
37	284.4
38	284.4
39	284.4
40	284.4
41	284.4
42	284.4
43	284.4
44	284.4
45	284.4
46	284.4
47	284.4
48	284.4
49	284.4
50	284.4
51	284.4
52	284.4
53	284.4
54	284.4
55	284.4
56	284.4
57	284.4
58	284.4
59	284.4
60	284.4
61	284.4
62	284.4
63	284.4
64	284.4
65	284.4
66	284.4
67	284.4
68	284.4
69	284.4
70	284.4
71	284.4
72	284.4
73	284.4
74	284.4
75	284.4
76	284.4
77	284.4
78	284.4
79	284.4
80	284.4
81	284.4
82	284.4
83	284.4
84	284.4
85	284.4
86	284.4
87	284.4
88	284.4
89	284.4
90	284.4
91	284.4
92	284.4
93	284.4
94	284.4
95	284.4
96	284.4
97	284.4
98	284.4
99	284.4
100	284.4